

MATH 4370/7370 – Linear Algebra and Matrix Analysis

Nonnegative matrices

Julien Arino

Fall 2023



**University
of Manitoba**

Outline

Definitions and some preliminary results

The Perron-Frobenius theorem

Stochastic matrices

Definitions and some preliminary results

Zero-nonzero structure of a matrix

The Perron-Frobenius theorem

Stochastic matrices

Definition 6.1 (Nonnegative/positive matrix)

A matrix $A \in \mathcal{M}_{mn}(\mathbb{R})$ is a **nonnegative matrix** if $a_{ij} \geq 0$ for all $i = 1, \dots, m$ and $j = 1, \dots, n$. We write $A \geq 0$. A is a **positive matrix** if $a_{ij} > 0$ for all $i = 1, \dots, m$ and $j = 1, \dots, n$. We write $A > 0$

Remark 6.2

In other references, you will see

▶ $A \geq 0 \iff a_{ij} \geq 0$

▶ $A > 0 \iff A \geq 0$ and there exists (i, j) , $a_{ij} > 0$ *[positive]*

▶ $A \gg 0 \iff a_{ij} > 0$ for all i, j *[strongly positive]*

I tend to favour the latter notation over the one used in these notes, but since the former is more common in matrix theory, I use the notation of Definition 6.1 here

Notation

Let $A, B \in \mathcal{M}_{mn}(\mathbb{R})$. Nonnegativity and positivity are used to define partial orders on $\mathcal{M}_{mn}(\mathbb{R})$

$$\blacktriangleright A \geq B \iff A - B \geq 0$$

$$\blacktriangleright A > B \iff A - B > 0$$

The same is used for vectors $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$: $\mathbf{x} \geq \mathbf{y}$ and $\mathbf{x} > \mathbf{y}$ if, respectively, $\mathbf{x} - \mathbf{y} \geq \mathbf{0}$ and $\mathbf{x} - \mathbf{y} > \mathbf{0}$. Note that the order is only partial: if $A \geq 0$ and $B \geq 0$, for instance, it is not necessarily possible to decide on the ordering of A and B with respect to one another

Theorem 6.3

Let A and B be nonnegative matrices of appropriate sizes. Then $A + B$ and AB are nonnegative. If $A > 0$ and $B \geq 0$, $B \neq 0$, then $AB \geq 0$ and $AB \neq 0$

Corollary 6.4

Let $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ be such that $\mathbf{x} \geq \mathbf{y}$ and $A \in \mathcal{M}_{mn}$ be nonnegative. Then $A\mathbf{x} \geq A\mathbf{y}$. Assume additionally that $\mathbf{x} \geq \mathbf{y}$, $\mathbf{x} \neq \mathbf{y}$ and $A > 0$. Then $A\mathbf{x} > A\mathbf{y}$

Definitions and some preliminary results

Zero-nonzero structure of a matrix

The Perron-Frobenius theorem

Stochastic matrices

Definition 6.5

Let $P, Q \in \mathcal{M}_{nm}(\mathbb{F})$. P and Q have the same **zero-nonzero structure** if for all i, j ,
 $p_{ij} \neq 0 \iff q_{ij} \neq 0$

Zero-nonzero structure defines an equivalence relation. Therefore, as with all equivalence relations, one only needs one representative from the equivalence class. One typical representative is defined using Boolean matrices

Definition 6.6

A **Boolean matrix** is a matrix whose entries are Boolean $\{0, 1\}$ and use Boolean arithmetics:

- ▶ $0 + 0 = 0$
- ▶ $1 + 0 = 0 + 1 = 1$
- ▶ $1 + 1 = 1$
- ▶ $0 \cdot 1 = 1$ and $1 \cdot 0 = 0 = 0 \cdot 0$
- ▶ $1 \cdot 1 = 1$

Definition 6.7

Let $A \in \mathcal{M}_{nm}(\mathbb{F})$. Then A_B denotes the **Boolean representation** of A , defined as follows. If $A = [a_{ij}]$, then $A_B = [\alpha_{ij}]$ with

$$\alpha_{ij} = \begin{cases} 1 & \text{if } a_{ij} \neq 0 \\ 0 & \text{if } a_{ij} = 0 \end{cases}$$

Definitions and some preliminary results

The Perron-Frobenius theorem

The Perron-Frobenius Theorem for irreducible matrices

Proof of a result of Perron for positive matrices

Proof of the Perron-Frobenius theorem for irreducible matrices

Primitive matrices

The Perron-Frobenius Theorem for nonnegative matrices

The Perron-Frobenius Theorem (revamped)

Application of the Perron-Frobenius Theorem

Stochastic matrices

Definitions and some preliminary results

The Perron-Frobenius theorem

The Perron-Frobenius Theorem for irreducible matrices

Proof of a result of Perron for positive matrices

Proof of the Perron-Frobenius theorem for irreducible matrices

Primitive matrices

The Perron-Frobenius Theorem for nonnegative matrices

The Perron-Frobenius Theorem (revamped)

Application of the Perron-Frobenius Theorem

Stochastic matrices

Theorem 6.15 (Perron-Frobenius)

Let $A \geq 0 \in \mathcal{M}_n$ be irreducible. Then the spectral radius $\rho(A) = \max\{|\lambda|, \lambda \in \sigma(A)\}$ is an eigenvalue of A . It is simple (has algebraic multiplicity 1), positive and is associated with a positive eigenvector. Furthermore, there is no nonnegative eigenvector associated to any other eigenvalue of A

Remark 6.16

*We often say that $\rho(A)$ is the **Perron root** of A ; the corresponding eigenvector is the **Perron vector** of A*

Definitions and some preliminary results

The Perron-Frobenius theorem

The Perron-Frobenius Theorem for irreducible matrices

Proof of a result of Perron for positive matrices

Proof of the Perron-Frobenius theorem for irreducible matrices

Primitive matrices

The Perron-Frobenius Theorem for nonnegative matrices

The Perron-Frobenius Theorem (revamped)

Application of the Perron-Frobenius Theorem

Stochastic matrices

Lemma 6.17 (Perron)

Let $M_n \ni A > 0$. Then $\rho(A)$ is a positive eigenvalue of A and there is only one linearly independent eigenvector associated to $\rho(A)$, which can be taken to be positive

Lemma 6.18

Let $\alpha_1, \dots, \alpha_n \in \mathbb{R}_+^*$ and $v_1, \dots, v_n \in \mathbb{C}$. Then

$$\left| \sum_{i=1}^n \alpha_i v_i \right| \leq \sum_{i=1}^n \alpha_i |v_i| \quad (1)$$

with equality if and only if there exists $\eta \in \mathbb{C}$, $|\eta| = 1$, such that $\eta v_i \geq 0$ for all $i = 1, \dots, n$

Definitions and some preliminary results

The Perron-Frobenius theorem

The Perron-Frobenius Theorem for irreducible matrices

Proof of a result of Perron for positive matrices

Proof of the Perron-Frobenius theorem for irreducible matrices

Primitive matrices

The Perron-Frobenius Theorem for nonnegative matrices

The Perron-Frobenius Theorem (revamped)

Application of the Perron-Frobenius Theorem

Stochastic matrices

Theorem 6.19

Let $A \in \mathcal{M}_n$ and $f(x)$ a polynomial. Then

$$\sigma(f(A)) = \{f(\lambda_1), \dots, f(\lambda_n), \lambda_i \in \sigma(A)\}$$

If we have $g(\lambda_i) \neq 0$ for $\lambda_i \in \sigma(A)$, for some polynomial g , then the matrix $g(A)$ is non-singular and

$$\sigma(f(A)g(A)^{-1}) = \left\{ \frac{f(\lambda_1)}{g(\lambda_1)}, \dots, \frac{f(\lambda_n)}{g(\lambda_n)}, \lambda_i \in \sigma(A) \right\}$$

If $x \neq 0$ eigenvector of A associated to $\lambda \in \sigma(A)$, then x is also an eigenvector of $f(A)$ and $f(A)g(A)^{-1}$ associated to eigenvalue $f(\lambda)$ and $f(\lambda)/g(\lambda)$, respectively

Lemma 6.20 (Schur's lemma)

Let $A \in \mathcal{M}_n$ and $\lambda \in \sigma(A)$. Then λ is simple if and only if both the following conditions are satisfied:

1. There exists only one linear independent eigenvector of A associated to λ , say \mathbf{u} , and thus only one linear independent eigenvector of A^T associated to λ , say \mathbf{v}
2. Vectors \mathbf{u} and \mathbf{v} in (1) satisfy $\mathbf{v}^T \mathbf{u} \neq 0$

Definitions and some preliminary results

The Perron-Frobenius theorem

The Perron-Frobenius Theorem for irreducible matrices

Proof of a result of Perron for positive matrices

Proof of the Perron-Frobenius theorem for irreducible matrices

Primitive matrices

The Perron-Frobenius Theorem for nonnegative matrices

The Perron-Frobenius Theorem (revamped)

Application of the Perron-Frobenius Theorem

Stochastic matrices

Definition 6.21

Let $\mathcal{M}_n(\mathbb{R}) \ni A \geq 0$. We say that A is **primitive** (with **primitivity index** $k \in \mathbb{N}_+^*$) if there exists $k \in \mathbb{N}_+^*$ such that

$$A^k > 0$$

with k the smallest integer for which this is true. We say that a matrix is **imprimitive** if it is not primitive

Remark 6.22

Primitivity implies irreducibility. The converse is not true

Theorem 6.23

A sufficient condition for primitivity is irreducibility with at least one positive diagonal entry

Here d is the index of imprimitivity (i.e., the number of eigenvalues that have the same modulus as $\lambda_p = \rho(A)$). If $d = 1$, then A is primitive. We have that $d = \text{gcd}$ of all the lengths of closed walks in $G(A)$

Theorem 6.24

Let $A \in \mathcal{M}_n$ be a non-negative matrix. If A is primitive, then $A^k > 0$ for some $0 < k \leq (n-1)n^n$

Theorem 6.25

Let $A \geq 0$ primitive. Suppose the shortest simple directed cycle in $G(A)$ has length s , then primitivity index is $\leq n + s(n - 1)$

Theorem 6.26

Let $A \in \mathcal{M}_n$ be a nonnegative matrix. A is primitive if and only if $A^{n^2-2n+2} > 0$

Theorem 6.27

Let $A \in \mathcal{M}_n$ be a nonnegative irreducible matrix. Suppose that A has d positive entries on the diagonal. Then the primitivity index is $\leq 2n - d - 1$

Definitions and some preliminary results

The Perron-Frobenius theorem

The Perron-Frobenius Theorem for irreducible matrices

Proof of a result of Perron for positive matrices

Proof of the Perron-Frobenius theorem for irreducible matrices

Primitive matrices

The Perron-Frobenius Theorem for nonnegative matrices

The Perron-Frobenius Theorem (revamped)

Application of the Perron-Frobenius Theorem

Stochastic matrices

Theorem 6.28

Let $A \geq 0$ in \mathcal{M}_n . Then there exists $0 \neq v \geq 0$ such that $Av = \rho(A)v$

Definitions and some preliminary results

The Perron-Frobenius theorem

The Perron-Frobenius Theorem for irreducible matrices

Proof of a result of Perron for positive matrices

Proof of the Perron-Frobenius theorem for irreducible matrices

Primitive matrices

The Perron-Frobenius Theorem for nonnegative matrices

The Perron-Frobenius Theorem (revamped)


Application of the Perron-Frobenius Theorem

Stochastic matrices

Let us restate the Perron-Frobenius theorem, taking into account the different cases. The classification in the following result is inspired by the presentation in [?].

Theorem 6.29

Let $\mathcal{M}_n \ni A \geq 0$. Denote λ_P the Perron root of A , i.e., $\lambda_P = \rho(A)$, \mathbf{v}_P and \mathbf{w}_P the corresponding right and left Perron vectors of A , respectively. Denote d the index of imprimitivity of A (with $d = 1$ when A is primitive) and $\lambda_j \in \sigma(A)$ the spectrum of A , with $j = 2, \dots, n$ unless otherwise specified (assuming $\lambda_1 = \lambda_P$). Then conclusions of the Perron-Frobenius Theorem can be summarised as follows.



Definitions and some preliminary results

The Perron-Frobenius theorem

The Perron-Frobenius Theorem for irreducible matrices

Proof of a result of Perron for positive matrices

Proof of the Perron-Frobenius theorem for irreducible matrices

Primitive matrices

The Perron-Frobenius Theorem for nonnegative matrices

The Perron-Frobenius Theorem (revamped)

Application of the Perron-Frobenius Theorem

Stochastic matrices

Theorem 6.30

Let $A \in \mathcal{M}_n$ be a nonnegative irreducible matrix and $\rho(A) \in \mathbb{N}_+$. Then the following are equivalent:

1. there exists exactly h distinct eigenvalues such that $|\lambda| = \rho(A)$.
2. there exists P a permutation matrix such that




$$PAP^T = \begin{pmatrix} 0 & A_{12} & 0 & \dots & 0 \\ \vdots & & A_{23} & \ddots & \vdots \\ \vdots & & & \ddots & 0 \\ \vdots & & & & A_{h-1,h} \\ A_{h1} & 0 & \dots & \dots & 0 \end{pmatrix}$$

where the diagonal blocks are square, and there does not exist other permutation matrix giving less than h horizontal blocks.

3. the greatest common divisor of the lengths of all cycles in $G(A)$ is h .
4. h is the maximal positive integer k such that

Corollary 6.31

Let $A \in \mathcal{M}_n$, $A \geq 0$ irreducible with exactly h distinct eigenvalues of modulus $\rho(A)$. Then, we can consider these eigenvalues as points in the complex plan, the eigenvalues are the vertices of a regular polygon of h sides with centre at the origin and are of the vertices being $\rho(A)$



Remark 6.32

For Fiedler, a primitive matrix is defined as an irreducible nonnegative matrix such that $h = 1$

Theorem 6.33

Let $A \geq 0$ in \mathcal{M}_n , $n \geq 2$. TFAE

1. $A^n = 0$
2. there exists $\mathbb{N} \ni k > 0$ such that $A^k = 0$
3. $G(A)$ acyclic
4. $\exists P$, permutation matrix, .t. PAP^T is upper-triangular with zeros on main diagonal
5. $\rho(A) = 0$

Theorem 6.34

Let $A \geq 0$ be a nonnegative matrix in \mathcal{M}_n . Assume that A has a positive eigenvector. Then that eigenvector is the Perron vector and is associated to $\rho(A)$

Definitions and some preliminary results

The Perron-Frobenius theorem

Stochastic matrices

- Row- and column-stochastic matrices

- Doubly stochastic matrices

Definitions and some preliminary results

The Perron-Frobenius theorem

Stochastic matrices

Row- and column-stochastic matrices

Doubly stochastic matrices

Definition 6.35 (Stochastic matrix)

The matrix $A \in \mathcal{M}_n$ is **stochastic** if

- $A \geq 0$ [The matrix is nonnegative]
- $A\mathbb{1} = \mathbb{1}$, $\mathbb{1} = (1, \dots, 1)^T$ [All rows sum to 1]

Equivalently, the matrix is stochastic if its column sums all equal 1

Definition 6.36

The matrix is **row-stochastic** or **column-stochastic**, respectively, if the rows or columns sum to 1. The terms **right stochastic** and **left stochastic** are also used. If both rows and columns sum to 1, then the matrix is **doubly stochastic**

Theorem 6.37

Let $A \in \mathcal{M}_n$ be stochastic. Then $\rho(A) = 1$

Theorem 6.38

Let $P \in \mathcal{M}_n$, $P \geq 0$. Assume that P has a positive eigenvector u and that $\rho(P) > 0$. Then there exists D , diagonal matrix with $\text{diag}(D) > 0$, and $k > 0$, $k \in \mathbb{R}$ such that

$$A = kDPD^{-1}$$

is stochastic, with $k = \rho(P)^{-1}$

Theorem 6.39

Let $A, B \in \mathcal{M}_n$ be stochastic. Then AB is stochastic

Theorem 6.40

Let A be a primitive stochastic. Then $A^k \rightarrow \mathbb{1}\mathbf{v}^T$, $k \rightarrow \infty$, where $\mathbb{1}\mathbf{v}^T$ has rank 1 and \mathbf{v} is the (left) eigenvector of A^T associated to $\rho(A) = 1$ and normalised so that $\mathbf{v}^T \mathbb{1} = 1$

Remark 6.41

This is a result that is used to compute the limit of a regular Markov chain

Definitions and some preliminary results

The Perron-Frobenius theorem

Stochastic matrices

Row- and column-stochastic matrices

Doubly stochastic matrices

Definition 6.42

The matrix $A \in \mathcal{M}_n$, $A \geq 0$ is **doubly stochastic** if $A\mathbb{1} = \mathbb{1}$ and $\mathbb{1}^T A = \mathbb{1}^T$

Remark 6.43

Here $\rho(A) = 1$ is associated to $\mathbb{1}$ for A and for A^T

Consider E the Euclidean space. A set K of points in E is **convex** if A_1, A_2 points in K , $\lambda_1, \lambda_2 \in \mathbb{R}_+$ such that $\lambda_1 + \lambda_2 = 1$, then

$$\lambda_1 A_1 + \lambda_2 A_2 \in K.$$

A **convex polyhedron** K is the set of all points of the form

$$\sum_{i=1}^N \lambda_i A_i$$

where A_i are points in E and $\lambda_i \in \mathbb{R}_+$

Let $A = [a_{ij}] \in \mathcal{M}_n(\mathbb{R})$. Consider this matrix as a point in E with coordinates $[a_{11}, a_{12}, \dots, a_{nn}]$ ($\dim E = n^2$)


Theorem 6.44

Let $A \in \mathcal{M}_n$, $A = [a_{ij}]$, if A is doubly stochastic, then this forms an $(n - 1)^2$ dimensional subspace of $\tilde{E} = \mathbb{R}^{n^2}$

Theorem 6.45 (Birkhoff)

In the space $\tilde{E} = \mathbb{R}^{n^2}$, the set of doubly stochastic matrices of order n is a convex polyhedron in E (the subspace of stochastic matrices). The vertices of the polyhedron are the points corresponding to all the permutation matrices

References I

-  Miroslav Fiedler, *Special matrices and their applications in numerical mathematics*, Dover, 2008.